

AD-A286 214



ATION PAGE

Form Approved
OMB No. 0704-0188

1. Purpose: This form is to be used for recording the time for reviewing instructions, searching existing data sources, gathering the collection of information, and comments regarding this burden estimate or any other aspect of this form, including suggestions for reducing the burden. Send comments to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302.

2. REPORT DATE October 18, 1994		3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE A Catalytic Method for the Conversion of Silanes to Stannanes		5. FUNDING NUMBERS Grant #N00014-91-J-1043 R&T Code 3135014 Harold E. Guard, Program Officer	
6. AUTHOR(S) Benjamin P. Warner and Stephen L. Buchwald		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Massachusetts Institute of Technology Department of Chemistry Cambridge, MA 02139		10. SPONSORING / MONITORING AGENCY REPORT NUMBER Technical Report No. 5	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Department of the Navy Chemistry Division, ONR 331 800 North Quincy Street Arlington, VA 22217-5660		11. SUPPLEMENTARY NOTES Prepared for Publication in the Journal of Organic Chemistry	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release and sale, distribution unlimited.		12b. DISTRIBUTION CODE 94-35065 	
13. ABSTRACT (Maximum 200 words) The transformation of alkynyl-, allyl-, and benzyltrimethylsilanes to the corresponding tributylstannanes is reported. The reaction is initiated by the addition of tetrabutylammonium fluoride to a mixture of the silane and bis(tributyltin)oxide in tetrahydrofuran. The stannanes are isolated in ~quantitative yields after removal of the volatile bis(trimethylsilyl)oxide <i>in vacuo</i> .			
14. SUBJECT TERMS silane, stannane, catalysis		15. NUMBER OF PAGES 9	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	
19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified		20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

OFFICE OF NAVAL RESEARCH

GRANT #N00014-91-J-1043

**R&T Code 3135014
Harold E. Guard, Program Officer
Chemistry Division**

Technical Report No. 5

**A Catalytic Method for the Conversion
of Silanes to Stannanes**

by

Benjamin P. Warner and Stephen L. Buchwald

Prepared for Publication

in the

Journal of Organic Chemistry

**Massachusetts Institute of Technology
Department of Chemistry
Cambridge, MA 02139**

October 18, 1994

**Reproduction in whole or in part is permitted for any purpose of the United States
Government**

A Catalytic Method for the Conversion of Silanes to Stannanes

Benjamin P. Warner and Stephen L. Buchwald*

Department of Chemistry

Massachusetts Institute of Technology

Cambridge, Massachusetts 02139

ABSTRACT


The transformation of alkynyl-, allyl-, and benzyltrimethylsilanes to the corresponding tributylstannanes is reported. The reaction is initiated by the addition of tetrabutylammonium fluoride to a mixture of the silane and bis(tributyltin)oxide in tetrahydrofuran. The stannanes are isolated in -quantitative yields after removal of the volatile bis(trimethylsilyl)oxide *in vacuo*.

Revised

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

In the course of our studies of organo-main group compounds, we desired ready access to aryl(alkynyl)boranes. One of the most useful routes to such compounds was from the aryl(silyl)acetylenes generated by the palladium mediated coupling of an aryl R-SiMe_3 and trimethylsilylacetylene.¹ The trimethylsilyl group would then be converted to a tributylstannyl group, and then to the corresponding alkynylborane.² Rather than the usual two step methods for converting alkynylsilanes to alkynylstannes,³ we reasoned that the trimethylsilyl group could be removed under appropriate conditions, and the resulting anion would react with bis(tributyltin)oxide. This reaction would generate a new alkoxide, and the cycle would be repeated. We found that tetrabutylammonium fluoride (TBAF) is an excellent catalyst for the process, the presumed course of which is shown in Scheme

Scheme 1



We believe that this method for generating alkynylstannanes from silanes has advantages in terms of cost and ease of use. This reaction utilizes inexpensive bis(tributyltin)oxide rather than the more costly and moisture-sensitive tributyltin chloride.⁵ The reaction allows the conversion of alkynylsilanes, as well as allyl- and benzyltrimethylsilane, to the corresponding tributylstannanes in one step, as opposed to desilylation and isolation of the terminal alkyne.³ Finally, the product is isolated in

quantitative yield with removal of the volatile bis(trimethylsilyl)oxide the only purification needed.⁶

The reaction was carried out by first charging a sealable Schlenk tube with an appropriate silane (1 to 1.05⁷ equiv), bis(tributyltin)oxide (0.5 equiv), and THF. A small amount of TBAF (0.02 equiv) was then added and the solution was heated at 60 °C for 2.5 h (16 h for allyl and benzyl silane), at which time the solvent and bis(trimethylsilyl)oxide are removed *in vacuo*. Allyl,⁸ benzyl, and alkynylsilanes all react to generate the corresponding stannanes in excellent yields without further purification (Table 1).

[Table 1]

In summary, we have developed an efficient method for the conversion of alkynyl, benzyl, and allyl silanes to the corresponding stannanes.

Experimental Section:

General Considerations:

All reactions were carried out under an atmosphere of argon using standard Schlenk techniques. Nuclear magnetic resonance (NMR) spectra were recorded on a Varian Unity-300, Varian XL-300, Varian XL 301, or Bruker AC-250 Fourier transform spectrometer. Infrared (IR) spectra were recorded on a Perkin Elmer 1600 series Fourier transform spectrometer. Electron impact high resolution mass determinations (HRMS) were recorded on a Finnegan MAT System 8200. Elemental Analyses were performed by Desert Analytics; air sensitive samples were sent in sealed vials under nitrogen.

Tetrahydrofuran (THF) was dried and deoxygenated by refluxing and distilling from sodium / benzophenone ketyl under an argon atmosphere. All reagents, unless otherwise stated, are commercially available and were used as received. Yields refer to isolated yields of products of greater than 95% purity as estimated by ¹H NMR

spectrometry, and are an average of two or more separate experiments.

Representative Procedures: A flame dried sealable Schlenk flask under Argon was charged with **1a** (0.348 g, 2.0 mmol), and $(\text{Bu}_3\text{Sn})_2\text{O}$ (0.596 g, 1.0 mmol), and THF (5 mL). TBAF (0.040 mL, 1 M in THF) was added, and the flask was sealed and stirred at 60 °C for 2.5 h, at which time the volatiles were removed *in vacuo* to yield **1b** as a colorless oil with no further purification necessary (0.764 g, 98 %).

Compounds **1a-4a**, **9a**, and **10a** were purchased from Aldrich Chemical Co., Inc. Compounds **5a-8a** were prepared according to the literature. The spectral data for **6a**,⁹ **1b**,¹⁰ and **2b**,¹¹ have been reported in the literature. Compounds **3b** and **9b** were compared with material purchased from Aldrich Chemical Co., Inc.

5a:¹ ^1H NMR (300 MHz, CDCl_3): δ 0.26 (s, 9H), 7.54 (d, $J = 18.0$ Hz, 2H), 7.57 (d, $J = 18.0$ Hz, 2H); IR (Film): alkyne 2158, nitrile 2254.

7a:¹² ^1H NMR (300 MHz, CDCl_3): δ 0.18 (s, 9H), 1.59 (m, 4H), 2.11 (m, 4H), 6.18 (m, 1H); IR (Film): alkyne 2146.

8a:¹ ^1H NMR (300 MHz, CDCl_3): δ 0.34 (s, 18H), 7.28 (dd, $J = 5.7, 3.3$ Hz, 2H), 7.51 (dd, $J = 5.7, 3.3$ Hz, 2H); IR (Film): alkyne 2161.

4b:¹³ ^1H NMR (300 MHz, CDCl_3): δ 0.89 (t, $J = 7.5$ Hz, 9H), 1.00 (t, $J = 8.3$ Hz, 6H), 1.30 (m, 6H), 1.55 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3): δ 92.91, 83.94, 28.79, 26.97, 13.57, 11.36; IR (Film): alkyne 2036.

5b: ^1H NMR (300 MHz, CDCl_3): δ 0.91 (t, $J = 7.3$ Hz, 9H), 1.07 (t, $J = 8.0$ Hz, 6H), 1.36 (m, 6H), 1.6 (m, 6H), 7.51 (d, $J = 8.2$ Hz, 2 H), 7.53 (d, $J = 8.2$ Hz, 2 H); ^{13}C NMR (75 MHz, CDCl_3): δ 132.2, 131.7, 128.7, 118.5, 110.5, 108.0, 99.9, 28.9, 27.0, 13.7, 11.3; IR (Film): nitrile 2363, alkyne 2228; HRMS: Calcd. for $\text{C}_{21}\text{H}_{31}\text{NSn}$: 417.1478. Found: 417.1476.

6b:¹⁴ ^1H NMR (300 MHz, CDCl_3): δ 0.88 (t, $J = 7.5$ Hz, 9H), 0.99 (t, $J = 8.1$ Hz, 6H), 1.32 (m, 6H), 1.53 (m, 6H), 3.37 (s, 3H), 4.09 (s, 2H); IR (Film): alkyne 2149.

7b:¹⁵ ¹H NMR (300 MHz, CDCl₃): δ 0.89 (t, *J* = 7.2 Hz, 9H), 0.98 (t, *J* = 8.1 Hz, 6H), 1.34 (m, 10H), 1.57 (m, 10H), 2.09 (m, 4H), 6.10 (m, 1H); IR (Film): alkyne 2126.

8b: ¹H NMR (300 MHz, CDCl₃): δ 0.92 (t, *J* = 7.2 Hz, 18H), 1.06 (t, *J* = 12 Hz, 12H), 1.36 (m, 12H), 1.6 (m, 12H), 7.17 (dd, *J* = 5.7, 3.4 Hz, 2 H), 7.42 (dd, *J* = 5.8, 3.4 Hz, 2 H); ¹³C NMR (75 MHz, CDCl₃): δ 132.6, 127.2, 126.3, 108.5, 97.4, 28.9, 27.0, 13.6, 11.2; IR (Film): alkyne 2135; Anal: Calcd. for: C₃₄H₅₈Sn₂: C, 57.99; H, 8.3. Found: C, 57.97; H, 8.58.

10b:¹⁶ ¹H NMR (300 MHz, CDCl₃): δ 0.81 (t, *J* = 7.8 Hz, 6H), 0.87 (t, *J* = 7.4 Hz, 9H), 1.28 (m, 6H), 1.4 (m, 6H), 2.3 (t, *J*_{Sn-H} = 27 Hz, 2H), 6.9 (m, 3 H), 7.16 (m., 2 H).

Acknowledgment: This work was supported, in part, by the Office of Naval Research to whom we are grateful. SLB acknowledges additional support as a Camille & Henry Dreyfus Teacher-Scholar.

Supplementary material available: Copies of ¹H and ¹³C NMR spectrum for **5b** (2 pages).

References:

1. Takahashi, S.; Kuroyama, Y.; Sonogashira, K.; Hagihara, N. *Synthesis*, **1980**, 627.
2. Wrackmeyer, B.; Nöth, H. *Chem. Ber.* **1977**, *110*, 1086.
3. Cf. Williamson, B.L.; Tykwynski, R.R.; Stang, P.J. *J. Am. Chem. Soc.* **1994**, *116*, 93. and references therein.
4. Nakamura, E.; Kuwajima, I. *Angew. Chem. Int. Ed. Eng.* **1976**, *15*, 498.
5. Tributyltin chloride is \$47.59/mole and bis(tributyltin)oxide is \$33.65/mole of tributyltin from Aldrich Chemical Co., Inc. Prices are from the 1992-93 catalog, and are derived from the largest sized bottles listed.
6. We note that the stannanes generated can be used without purification or removal of

solvent and volatiles, in Stille reactions. For example, **1a** was converted to **1b**, then 1 equivalent of iodobenzene, 0.025 equivalents of palladium (II) acetate, and 0.05 equivalents of triphenylphosphine were added. After heating the reaction mixture to 60 °C for 14 h, followed by aqueous work up, and flash chromatography, diphenylacetylene was isolated in 81 % yield. Echavarren, A.M.; Stille, J.K. *J. Am. Chem. Soc.* **1987**, *109*, 5478. Cummins, C.H. *Tetrahedron Lett.* **1994**, *35*, 857.

7. Solutions of TBAF contain 5% water, which protonates some of the silane or stannane. For alkynylstannanes, this reaction is reversible, but for allyl and benzyl trimethylsilane, a small amount of material was unavoidably lost.

8. Hosomi, A.; Shirahata, A.; Sakurai, H. *Tetrahedron Lett.* **1978**, 3043.

9. Labaudinière, L.; Hanaizi, J.; Normant, J.F. *J. Org. Chem.* **1992**, *57*, 6903.

10. Loque, M.W.; Teng, K. *J. Org. Chem.* **1982**, *47*, 2549.

11. Keinan, E.; Peretz, M. *J. Org. Chem.* **1983**, *48*, 5302.

12. Stadnichuk, M.D.; Petrov, A.A. *Zh. Obschch. Khim.* **1961**, *31*, 411. [Chem. Abstr. **1961**, *55*, 23329a.]

13. Hartmann, H.; Wagner, H.; Karbstein, B.; el A'ssar, M.K.; Reiss, W. *Naturwissenschaften*, **1964**, *51*, 215.

14. Zhai, D., Zhai, W.; Williams, R.M. *J. Am. Chem. Soc.* **1988**, *110*, 2501. Williamson, B.L.; Stang, P.J.; Arif, A.M. *J. Am. Chem. Soc.* **1993**, *115*, 2590.

15. Ito, Y.; Inouye, M.; Murakami, M. *Tetrahedron Lett.* **1988**, *29*, 5379.

16. Davies, A.G.; Roberts, B.P.; Smith, J.M. *J. Chem. Soc., Perkin Trans. 2* **1972**, 2221.

Table 1

Substrate ^a		Product	Yield ^c
1a	<chem>Ph-C#C-SiMe3</chem>	1b <chem>Ph-C#C-SnBu3</chem>	97%
2a	<chem>Me-C#C-SiMe3</chem>	2b <chem>Me-C#C-SnBu3</chem>	98%
3a	<chem>Me3Si-C#C-SiMe3</chem>	3b <chem>Bu3Sn-C#C-SnBu3</chem>	98%
4a	<chem>Me3Si-C#C-C#C-SiMe3</chem>	4b <chem>Bu3Sn-C#C-C#C-SnBu3</chem>	98%
5a	<chem>p-NC-Ph-C#C-SiMe3</chem>	5b <chem>p-NC-Ph-C#C-SnBu3</chem>	99%
6a	<chem>MeO-CH2-C#C-SiMe3</chem>	6b <chem>MeO-CH2-C#C-SnBu3</chem>	98%
7a	<chem>C1=CCCCC1-C#C-SiMe3</chem>	7b <chem>C1=CCCCC1-C#C-SnBu3</chem>	97%
8a	<chem>c1ccccc1C#CC#CSiMe3</chem>	8b <chem>c1ccccc1C#CC#CSnBu3</chem>	99%
9a ^b	<chem>CH2=CH-CH2-SiMe3</chem>	9b <chem>CH2=CH-CH2-SnBu3</chem>	95%
10a ^b	<chem>Ph-CH2-SiMe3</chem>	10b <chem>Ph-CH2-SnBu3</chem>	95%

^aReaction run for 2.5 h except as noted.^bReaction run for 16 h.^cYields refer to isolated product of >95% purity as estimated by ¹H NMR.

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

**Office of Naval Research (1)*
Chemistry Division, ONR 331
800 North Quincy Street
Arlington, Virginia 22217-5660**

**Dr. Richard W. Drisko (1)
Naval Facilities & Engineering
Service Center
Code L52
Port Hueneme, CA 93043**

**Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314**

**Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198**

**Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5320**

**Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000**

**Dr. John Fischer, Director (1)
Chemistry Division, C0235
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001**

**Dr. Peter Seligman (1)
Naval Command, Control and
Ocean Surveillance Center
RDT&E Division
San Diego, CA 92152-5000**

* Number of copies to forward